

Target diseases	Recommended dosage	Application window	Application conditions
Anthracnose (<i>Colletotrichum</i> spp.)	0,1 - 0,3 kg/ha	From the beginning of vegetation period every 7-14 days	Unlimited number of spray applications. Recommended to mix in at least 120 l/ha.
Grey Mould (<i>Botrytis cinerea</i>)	0,1 - 0,3 kg/ha	From the beginning of vegetation period every 7-14 days	Unlimited number of spray applications with special emphasis on applications from pre-flowering. Recommended to mix in at least 120 l/ha.
Red Stele Root Rot (<i>Phytophthora fragariae</i>) Phytophthora Crown Rot (<i>Phytophthora parasitica</i>)	0,05 %		Dipping of seedlings before planting, watering after planting
	0,1 - 0,3 kg/ha	After planting*	Unlimited number of spray/drip applications with special emphasis on early applications after planting. Recommended to mix in at least 120 l/ha.
Botryosphaeria stem blight (<i>Lasiodiplodium theobromae</i>)	0,3 kg/ha	After pruning	Preventive application. At least 2 applications after pruning timed 5–7 days apart. Recommended to mix in 120 l of water, but preferably more.
Common leaf spot (<i>Mycosphaerella fragariae</i>)	0,1 - 0,3 kg/ha	First application at the beginning of flowering. Start follow-up applications in late flowering every 7-14 days.	Unlimited number of spray applications. Adjust the number of sprays according to the expected infestation pressure. Recommended to mix in at least 120 l/ha.

* If seedlings are pre-treated, follow up with the first application 7-14 days after planting